

Local climate profile

Flinders Municipality



Past and current climate:

- The Flinders municipality includes all the islands of the Furneaux group in eastern Bass Strait. These islands have a maritime climate influenced by the surrounding seas and the mid-latitude westerly circulation (the 'Roaring 40s'). Temperatures are generally temperate with a small seasonal range (Flinders Island airport has an average daily maximum temperature of around 23 °C in February, and 13 °C in July).
- Flinders Island has a moderate rainfall of around 600 mm at the coast up to around 800 mm in the hills, with a moderate seasonal cycle. For example, Flinders Island airport receives 743 mm (39 mm in February, 84 mm in July) and Memana receives 779 mm (41 mm in February, 80 mm in July).
- Rainfall in the municipality comes from a variety of sources, including some from the regular westerly frontal rain systems that cross Tasmania. However, a significant proportion of the rainfall comes from episodic systems from the north and east, including cutoff lows.
- Year-to-year rainfall variability in this area shows a correlation with the El Niño Southern Oscillation in autumn, winter and spring (where El Niño winters are generally drier than average, La Niña winters are generally wetter than average). There is also a correlation with atmospheric blocking in almost all seasons (blocking affects the incidence of easterly systems and cutoff lows). The temperature and rainfall of the region is influenced by the warm waters brought south by the East Australia Current.
- Average temperatures have risen in the decades since the 1950s, at a rate similar to the rest of Tasmania (up to 0.15 °C per decade). Daily minimum temperatures have risen slightly more than daily maximum temperatures.
- There has been a decline in average rainfall and a lack of very wet years in the Flinders municipality since the mid 1970s, and this decline has been strongest in autumn. This decline was exacerbated by the 'big dry' drought of 1995-2009. The recent period since the end of the drought have seen rainfalls that are close to average or above.

Future scenarios - from the Climate Futures for Tasmania project

Fine-scale model projections of Tasmanian climate were made for two hypothetical but plausible scenarios of human emissions for the 21st Century (taken from the special report on emissions scenarios (SRES) from the Intergovernmental Panel on Climate Change (IPCC)). The scenarios are of ongoing high emissions, A2, and one where emissions plateau and fall, B1. The climate response under the two scenarios is similar through the first half of the century, but the changes under the higher emissions scenario become much stronger than the lower scenario in the later half of the 21st Century.

1. Temperature

- Under the higher emissions scenario (A2), the municipality is projected to experience a rise in average temperatures of 2.6 to 3.3 °C over the entire 21st Century. The rise in daily minimum temperature is expected to be slightly greater than daily maximum temperature, and fairly similar in the different seasons. Under the lower emissions scenario (B1), the projected change over the entire century is 1.3 to 2.0 °C. A time series of projected mean Tasmanian temperature is shown in Figure 1.
- The projected change in average temperatures is similar to the rest of Tasmania, but less than the global average and significantly less than northern Australia and many regions around the world, especially the large northern hemisphere continents and the Arctic.




Figure 1. Tasmanian average temperature in observations (black) and model projections for the A2 scenario (red) and the B1 scenario (blue), all series are smoothed (11-year running average), shading shows the range of model projections. Changes under the higher scenario by the very end of the century are discussed in the examples below.

- The projected change in average temperature is accompanied by a change in the frequency, intensity and duration of hot and cold extremes of temperature. For Flinders Island under the A2 (higher) scenario by the end of the century the projections indicate:
 - The number of Summer Days (>25 °C) increases from around 20 days per year, to more than 40 days per year. Nights with a minimum temperature >20 °C are projected to occur.
 - The temperature of very hot days increases more than the change in average temperature (by 3-4 °C in some locations in some seasons).
 - A reduction in frost-risk days to a very low frequency.
 - Warm spells (days in a row where temperatures are in the top 5% of baseline levels) currently last around 5 days, are projected to last up to 18 days longer.

2. Rainfall, runoff and rivers

- The projected pattern of change to rainfall and runoff is similar in nature between the two scenarios, but stronger by the end of the century under the A2 scenario. The model projections indicate that the general long-term influence of climate warming by the end of the century is for increased annual average rainfall at the coast.




Figure 2. Average rainfall in summer and winter – the left hand side plots show the average rainfall in the baseline period (1961-1990), the plots to the right show the proportional change (%) from that amount in various periods in the 21st century in the average of six climate model projections under the A2 (higher) emissions scenario.

- The model projections indicate an increase in annual average rainfall by the end of the century (up to 20% change by the end of the century under the higher emissions scenario).
- There is a projected increase in rainfall in all seasons, all generally in the 5-20% range by the end of the century under the higher emissions scenario. See Fig. 2 for summer and winter.
- The long-term effect of greenhouse warming is on top of the usual cycles of rainfall, including droughts, termed 'natural variability'. The model projections indicate that the recent dry conditions of the 'big dry' drought is not a new ongoing climate average state. These projections indicate that in the long term, drought frequency and severity on Flinders Island is actually projected to decrease due to the higher average rainfall.
- The projected changes to rainfall are driven by an strengthening of the East Australia Current bringing warmer waters to the area, as well as a change to the changes to the average circulation over the area and the incidence of rain-bearing systems from the east and north including cutoff lows.
- A major influence of greenhouse warming on rainfall is the tendency for heavier rainfalls interspersed by longer dry periods, and for greater extremes. However, this varies in different areas. For Flinders Island under the A2 (higher) scenario by the end of the century there is projected to be:

- Around 3 fewer days with >1 mm rain per year on average, but significantly more rain per rain day.
 - Up to 7 more very wet days each year (where rainfall exceeds the baseline 95th percentile).
 - An increase in the maximum instantaneous rainfall rate of over 35% in some seasons, and an increase of >15 mm of rainfall on the average wettest day of the year (up to 30% increase).
 - Rainfall brought by rare extreme events is projected to increase: a 200-year average recurrence interval (ARI) event for daily rainfall at is projected to increase by more than 50%. More common ARI events (ARI-10, ARI-50) are projected to increase by a similar proportion.
- Pan evaporation is projected to increase, by up to 19% under the A2 scenario by the end of the century, driven by the increases in temperature but also changes to relative humidity, wind speeds, cloudiness and radiation.
 - Changes to rainfall and evaporation lead to changes in water runoff and river flows. This in turn has impacts on the inflows into dams and water storages. Under the A2 scenario by the end of the century:
 - Average annual runoff amounts are projected to increase by a large amount (up to 60%) due to the increase in rainfall, with an increase in all seasons.
 - There is projected to be an increase in runoff during high events, also of up to 60% in some places. Runoff amounts during low events are projected to stay relatively similar to what they are now.
 - Projections of increased rainfall and runoff in all seasons result in large projected increase in average flows in Flinders Island rivers (central estimate is +46%).

3. Agricultural impacts

- Increasing temperatures are projected to increase the risk of problems from the Queensland fruit fly pest. The temperature regime on Flinders Island is projected to be suitable for permanent populations of fruit fly by the end of the century under the higher emissions scenario.
- The projected increase in rainfall leads to a corresponding reduction in the proportion of time spent in severe drought measured as the proportion of time with a standardized precipitation index (SPI) of less than minus two. This proportion is projected to fall from 3.3% to 0.9% under the higher emissions scenario by the end of the century.
- There is a projected increase in Growing Degree Days (GDD, a measure of the heat to grow and ripen crops). This will affect where crops can be grown, reduce the time to harvest of many crops, and affect many aspects of crop management.
- Frost risk days are projected to become much less frequent with a warming climate. Damaging spring frosts may still occur rarely.
- Chilling affects the growth and flowering of berries, fruits and nuts. Accumulated chill hours decrease given climate warming.
- For more information on agricultural impacts see Holz et al. (2011).

4. Extreme sea level events

High water events causing coastal inundation comes from a combination of sea level, tide, storm surge and wind waves. Sea level has been rising at a rate of 3.3 ± 0.4 mm/year in the recent period, and is expected to continue rising with further climate warming. The last IPCC assessment report gave a central estimate of a rise of 0.82 m global average sea level by 2100 under a high emissions scenario. The sea level rise varies in different locations, and for the coasts of Tasmania the sea level rise for this scenario is close to the global average.

The Furneaux group lies on the border between two different tidal regimes, with a higher tidal range on the north coast of Tasmania, and lower tidal ranges on the east coast. The difference in the influence of tide means there is a difference in the 1 in 100 year storm tide height: 1.4 to 1.5 m on the west coast of Flinders Island and 0.6 to 0.7 m on the east coast. Changes to storm surges by the end of the century will not be as large as sea level rise. Accounting for all effects, the 100-year event is projected to increase by the end of the century under the higher emissions scenario by an amount somewhat similar to Bicheno (an increase of just over 0.5 m).

Appendix – details of climate projections

Greenhouse gas emissions have an influence on the Earth's climate system, along with other human activities such as the emission of ozone-depleting substances, emission of aerosol (particles) and changing the land cover (e.g. deforestation). Sophisticated model simulations can be used to project the likely effect of these influences into the future given our current state of knowledge. It is impossible to predict exactly what future human emissions will be, so models are run under a set of plausible hypothetical emissions scenarios. A model simulation shows the likely effect if we follow that scenario, so it is not a single 'prediction' of the future. The simulation can't include the effect of things that are impossible to predict (such as major volcanic eruptions).

The Climate Futures for Tasmania project produced a set of climate projections at the regional scale for Tasmania. Two emissions scenarios were considered – one of ongoing high emissions (SRES A2), and one where emissions plateau and fall (SRES B1). The climate response under the two scenarios is similar through the first half of the century, but the changes under the higher emissions scenario become much stronger than the lower scenario in the latter half of the 21st Century.

Climate warming causes many complex changes to the earth's climate system. These changes include alterations to ocean currents, average atmospheric circulation and ocean-atmosphere cycles such as the El Niño Southern Oscillation. Projected effects that are relevant to Tasmania include a continued extension of the East Australia Current bringing warmer waters off the east and northeast coast of Tasmania, a pole-ward shift of the subtropical ridge of high pressure and shifts in the mid-latitude westerlies (the 'Roaring 40s'), and a change in remote climate drivers such as atmospheric blocking, the El Niño Southern Oscillation and the Southern Annular Mode. The position of Tasmania adjacent to the Southern Ocean means that the effect of climate warming is not as severe as other more continental regions.

The results presented in this report were made using established methods, including:

- Extreme value distribution fitting in a generalized Pareto distribution to calculate the average recurrence intervals (ARIs).
- Hydrology runoff models developed and calibrated for the Tasmanian Sustainable Yields project to estimate the runoff, river flows and inflows to storages.
- Standard agricultural indices such as the Utah

model to calculate chill hours and standard equations and a 10 °C threshold to calculate Growing Degree Days.

All information is drawn from the Climate Futures for Tasmania Technical reports please see these reports for more details, and to cite in other written work.

Reference list

- Bennett JC, Ling FLN, Graham B, Grose MR, Corney SP, White CJ, Holz GK, Post DA, Gaynor SM & Bindoff NL 2010, Climate Futures for Tasmania: water and catchments technical report, Antarctic Climate and Ecosystems Cooperative Research Centre, Hobart, Tasmania
- Corney SP, Katzfey JJ, McGregor JL, Grose MR, Bennett JC, White CJ, Holz GK, Gaynor SM & Bindoff NL 2011, Climate Futures for Tasmania: climate modeling technical report, Antarctic Climate and Ecosystems Cooperative Research Centre, Hobart, Tasmania
- Entura Consulting, 2010, Climate Futures for Tasmania Flood inundation mapping, Entura Consulting Technical report, 23 Dec 2010
- Grose MR, Barnes-Keoghan I, Corney SP, White CJ, Holz GK, Bennett JC, Gaynor SM & Bindoff NL 2010, Climate Futures for Tasmania: general climate impacts technical report, Antarctic Climate and Ecosystems Cooperative Research Centre, Hobart, Tasmania
- Holz GK, Grose MR, Bennett JC, Corney SP, White CJ, Phelan D, Potter K, Kriticos D, Rawnsley R, Parsons D, Lissos S, Gaynor SM & Bindoff NL 2010, Climate Futures for Tasmania: impacts on agriculture technical report, Antarctic Climate and Ecosystems Cooperative Research Centre, Hobart, Tasmania
- McInnes KL, O'Grady JG, Hemer M, Macadam I, Abbs DJ, White CJ, Bennett JC, Corney SP, Holz GK, Grose MR, Gaynor SM & Bindoff NL In Press, Climate Futures for Tasmania: extreme tide and sea level events technical report, Antarctic Climate and Ecosystems Cooperative Research Centre, Hobart, Tasmania
- White CJ, Sanabria LA, Grose MR, Corney SP, Bennett JC, Holz GK, McInnes KL, Cechet RP, Gaynor SM & Bindoff NL 2011, Climate Futures for Tasmania: extreme events technical report, Antarctic Climate and Ecosystems Cooperative Research Centre, Hobart, Tasmania

The material in this report is based on computer modelling projections for climate change scenarios and, as such, there are inherent uncertainties involved. While every effort has been made to ensure the material in this report is accurate, Antarctic Climate & Ecosystems Cooperative Research Centre (ACE) provides no warranty, guarantee or representation that material is accurate, complete, up to date, non-infringing or fit for a particular purpose. The use of the material is entirely at the risk of a user. The user must independently verify the suitability of the material for its own use.

To the maximum extent permitted by law, ACE, its participating organisations and their officers, employees, contractors and agents exclude liability for any loss, damage, costs or expenses whether direct, indirect, consequential including loss of profits, opportunity and third party claims that may be caused through the use of, reliance upon, or interpretation of the material in this report.