

## Local climate profile

# West Tamar Municipality



### Past and current climate:

- The West Tamar municipality has a temperate climate with moderate temperature range (near the coast the average daily maximum temperature is around 21 °C in February, and 12.5 °C in July).
- Average annual rainfall is generally 800-1000 mm per year with a distinct seasonal cycle. For example, Beauty Point receives 828 mm per year (40 mm in February, 102 in August), Beaconsfield receives 944 mm per year (46 mm in February and 121 mm in July) and Frankford receives 1059 mm per year (48 mm in February, 139 in July).
- Rainfall in the West Tamar area can come from the regular westerly frontal rain systems that cross Tasmania, however an important fraction of the rainfall comes from episodic systems from the north and east, including cutoff lows.
- Year-to-year rainfall variability is partly correlated with the El Niño Southern Oscillation in autumn, winter and spring (where El Niño winters are generally drier than average, La Niña winters are generally wetter than average). There is also some correlation with the Indian Ocean Dipole in winter and spring, and with atmospheric blocking especially in summer.
- Average temperatures have risen in the decades since the 1950s, at a rate similar to the rest of Tasmania (up to 0.15 °C per decade). Daily minimum temperatures have risen slightly more than daily maximum temperatures.
- There has been a general decline in average rainfall and a lack of very wet years in the West Tamar municipality since the mid 1970s, and this decline has been strongest in autumn. This decline was exacerbated by the 'big dry' drought of 1995-2009. The recent two years have seen above average rainfalls.


### Future scenarios - from the Climate Futures for Tasmania project

Fine-scale model projections of Tasmanian climate were made for two hypothetical but plausible scenarios of human emissions for the 21<sup>st</sup> Century (taken from the special report on emissions scenarios (SRES) from the Intergovernmental Panel on Climate Change (IPCC)). The scenarios are of ongoing high emissions, A2, and one where emissions plateau and fall, B1. The climate response under the two scenarios is similar through the first half of the century, but the changes under the higher emissions scenario become much stronger than the lower scenario in the later half of the 21<sup>st</sup> Century.

#### 1. Temperature

- Under the higher emissions scenario (A2), the municipality is projected to experience a rise in average temperatures of 2.6 to 3.3 °C over the entire 21<sup>st</sup> Century. The rise in daily minimum temperature is expected to be slightly greater than daily maximum temperature, and fairly similar in the different seasons. Under the lower emissions scenario (B1), the change over the entire century is projected to be 1.3 to 2.0 °C. A time series of projected mean Tasmanian mean temperature is shown in Fig 1.

- The projected change in average temperatures is similar to the rest of Tasmania, but less than the global average and significantly less than northern Australia and many regions around the world, especially the large northern hemisphere continents and the Arctic.




**Figure 1.** Tasmanian average temperature in observations (black) and model projections for the A2 scenario (red) and the B1 scenario (blue), all series are smoothed (11-year running average), shading shows the range of model projections. Changes under the higher scenario by the very end of the century are discussed in the examples below.

- The projected change in average temperature is accompanied by a change in the frequency, intensity and duration of hot and cold extremes of temperature. For West Tamar under the A2 (higher) scenario by the end of the century the projections indicate:
  - The number of Summer Days (>25 °C) increases from up to 20 days per year, to more than 45 days per year, with heat waves (3 days >28 °C) and night time minimum temperatures over 20 °C occurring occasionally.
  - The temperature of very hot days increases more than the change in average temperature (by 3-4 °C in some locations in some seasons).
  - A reduction in frost-risk days from up to 20 per year, to less than 5 per year.
  - Warm spells (days in a row where temperatures are in the top 5% of baseline levels) currently last around 7 days, are projected to last up to 14 days longer.

## 2. Rainfall, runoff and rivers

- The projected pattern of change to rainfall and runoff is similar in nature between the two scenarios, but stronger by the end of the century under the A2 scenario. The model projections indicate that the general long-term influence of climate warming by the end of the century is for an increase in annual average rainfall.



**Figure 2.** Average rainfall in summer and winter – the left hand side plots show the average rainfall in the baseline period (1961-1990), the plots to the right show the proportional change (%) from that amount in various periods in the 21<sup>st</sup> century in the average of six climate model projections under the A2 (higher) emissions scenario.

- There is a projected increase in annual average rainfall under the A2 scenario by the end of the century (model mean is for 0 to 10%). The tendency under the B1 scenario is for little change in annual rainfall.
- The model mean shows that rainfall is projected to remain much the same in autumn but increase in winter, spring and summer, with the strongest increase in winter (see Fig. 2 for summer and winter).
- The long-term effect of greenhouse warming is on top of the usual cycles of rainfall, including droughts, termed 'natural variability'. The model projections indicate that the recent dry conditions of the 'big dry' drought is not a new ongoing climate average state. These projections indicate that in the long term, drought frequency and severity may even reduce slightly due to the increase in average rainfall.
- The projected increase in rainfall is driven by changes to the average circulation of the region, including the average strength of the westerlies, as well as the incidence of the main rain-bearing weather systems from the east and north, including a change in atmospheric blocking and cutoff lows.

- A major influence of greenhouse warming on rainfall is the tendency for heavier rainfalls interspersed by longer dry periods, and for greater extremes. However, this varies in different areas. For the West Tamar municipality under the A2 (higher) scenario by the end of the century there is projected to be:
  - Up to 5 fewer days with >1 mm rain per year on average, but significantly more rain per rain day (a 15% increase or more).
  - Up to 3 more very wet days each year (where rainfall exceeds the baseline 95<sup>th</sup> percentile), and 2 more days per year that exceed 20 mm.
  - An increase in the maximum instantaneous rainfall rate of over 30% in some seasons, and an increase of 8 mm of rainfall on the wettest day of the year (a 20% increase).
  - An increase in the rainfall brought by rare extreme events: a 200-year average recurrence interval (ARI) event for daily rainfall at is projected to increase by more than 30 mm (a 35% increase). More common ARI events (ARI-10, ARI-50) are projected to increase by a similar proportion.
- Pan evaporation is projected to increase, by up to 19% under the A2 scenario by the end of the century, driven by the increases in temperature but also changes to relative humidity, wind speeds, cloudiness and radiation.
- Changes to rainfall and evaporation lead to changes in water runoff and river flows. This in turn has impacts on the inflows into dams and water storages. Under the A2 scenario by the end of the century:
  - Average runoff is projected to increase in all seasons.
  - Proportional (%) increases in average runoff are larger than the change to rainfall, with increases of over 20% possible in some seasons.
  - Runoff amounts during high events are projected to increase, including those events that may lead to erosion and localized flooding. Daily runoff amounts during low flows are projected to stay much the same or increase slightly.
  - Average flows in the Tamar River are projected to increase under the higher emissions scenario by the end of the century (central estimate is 19%).

### 3. Agricultural impacts

- Frost risk days are projected to become much less frequent with a warming climate. Damaging spring frosts may still occur rarely.
- Chilling affects the growth and flowering of berries, fruits and nuts. Accumulated chill hours decrease given the warming under the two future climate scenarios.
- Conditions to grow wine grapes are projected to change markedly, which will affect choices such as grape variety and harvest date. Vineyards in the West Tamar area currently experience around 1050 annual Biologically Effective GDDs, making them suitable for growing Pinot grapes. Under the A2 scenario, this is projected to change to around 1300 BEGDD in the coming decades, and up to 1800 BEGDD in the last decades of the century, making the temperature conditions similar to current conditions in the Coonawarra or even Rutherglen regions. If the vineyard continues growing Pinot grapes, the harvest date will move forward from June to February and the grape quality will be affected. Otherwise, the wine grower may plant other varieties of grapes, such as Shiraz and Cabernet Sauvignon. As well as temperature, other climate changes are projected to impact upon the wine industry, including heavy rain events leading to soil erosion and potential for fungal diseases.

- For more information on agricultural impacts, see Holz et al. (2011).

#### **4. Extreme sea level events**

High water events causing coastal inundation comes from a combination of sea level, tide, storm surge and wind waves. Sea level has been rising at a rate of  $3.3 \pm 0.4$  mm/year in the recent period, and is expected to continue rising with further climate warming. The last IPCC assessment report gave a central estimate of a rise of 0.82 m global average sea level by 2100 under a high emissions scenario. The sea level rise varies in different locations, and for the coasts of Tasmania the sea level rise for this scenario is close to the global average.

On the north coast of Tasmania, the very high tide height contributes more to coastal inundation events than the relatively modest very high storm surge height – the current 100-year storm tide event is around 1.96 m above average sea level. Changes to storm surges by the end of the century will not be as large as sea level rise. Accounting for all effects, the current 100-year event in George Town is projected to increase to 2.58 m by 2090 under the high emissions scenario.

#### **5. River floods – Tamar River**

Changes to design flood hydrographs were calculated for the 1:10, 1:50, 1:100 and 1:200 annual exceedance probability events for future periods using the climate model outputs and flood hydraulic models by partners at Entura consulting. Short duration high rainfall and runoff events are projected to become more intense, so catchments with a critical duration of less than 72 hours will experience high flood levels and faster response times. The peak discharge of the Tamar River is projected to increase significantly through the 21<sup>st</sup> Century under the higher emissions scenario. Additionally, sea level rise will significantly increase coastal inundation events in the estuary of the Tamar (see above point). Please see the full Entura report and accompanying maps for more details.

## Appendix – details of climate projections

Greenhouse gas emissions have an influence on the Earth's climate system, along with other human activities such as the emission of ozone-depleting substances, emission of aerosol (particles) and changing the land cover (e.g. deforestation). Sophisticated model simulations can be used to project the likely effect of these influences into the future given our current state of knowledge. It is impossible to predict exactly what future human emissions will be, so models are run under a set of plausible hypothetical emissions scenarios. A model simulation shows the likely effect if we follow that scenario, so it is not a single 'prediction' of the future. The simulation can't include the effect of things that are impossible to predict (such as major volcanic eruptions).

The Climate Futures for Tasmania project produced a set of climate projections at the regional scale for Tasmania. Two emissions scenarios were considered – one of ongoing high emissions (SRES A2), and one where emissions plateau and fall (SRES B1). The climate response under the two scenarios is similar through the first half of the century, but the changes under the higher emissions scenario become much stronger than the lower scenario in the latter half of the 21<sup>st</sup> Century.

Climate warming causes many complex changes to the earth's climate system. These changes include alterations to ocean currents, average atmospheric circulation and ocean-atmosphere cycles such as the El Niño Southern Oscillation. Projected effects that are relevant to Tasmania include a continued extension of the East Australia Current bringing warmer waters off the east and northeast coast of Tasmania, a pole-ward shift of the subtropical ridge of high pressure and shifts in the mid-latitude westerlies (the 'Roaring 40s'), and a change in remote climate drivers such as atmospheric blocking, the El Niño Southern Oscillation and the Southern Annular Mode. The position of Tasmania adjacent to the Southern Ocean means that the effect of climate warming is not as severe as other more continental regions.

The results presented in this report were made using established methods, including:

- Extreme value distribution fitting in a generalized Pareto distribution to calculate the average recurrence intervals (ARIs).
- Hydrology runoff models developed and calibrated for the Tasmanian Sustainable Yields project to estimate the runoff, river flows and inflows to storages.
- Standard agricultural indices such as the Utah model to calculate chill hours and standard

equations and a 10 °C threshold to calculate Growing Degree Days.

All information is drawn from the Climate Futures for Tasmania Technical reports please see these reports for more details, and to cite in other written work.

### Reference list

- Bennett JC, Ling FLN, Graham B, Grose MR, Corney SP, White CJ, Holz GK, Post DA, Gaynor SM & Bindoff NL 2010, Climate Futures for Tasmania: water and catchments technical report, Antarctic Climate and Ecosystems Cooperative Research Centre, Hobart, Tasmania
- Corney SP, Katzfey JJ, McGregor JL, Grose MR, Bennett JC, White CJ, Holz GK, Gaynor SM & Bindoff NL 2011, Climate Futures for Tasmania: climate modeling technical report, Antarctic Climate and Ecosystems Cooperative Research Centre, Hobart, Tasmania
- Entura Consulting, 2010, Climate Futures for Tasmania Flood inundation mapping, Entura Consulting Technical report, 23 Dec 2010
- Grose MR, Barnes-Keoghan I, Corney SP, White CJ, Holz GK, Bennett JC, Gaynor SM & Bindoff NL 2010, Climate Futures for Tasmania: general climate impacts technical report, Antarctic Climate and Ecosystems Cooperative Research Centre, Hobart, Tasmania
- Holz GK, Grose MR, Bennett JC, Corney SP, White CJ, Phelan D, Potter K, Kriticos D, Rawnsley R, Parsons D, Lissou S, Gaynor SM & Bindoff NL 2010, Climate Futures for Tasmania: impacts on agriculture technical report, Antarctic Climate and Ecosystems Cooperative Research Centre, Hobart, Tasmania
- McInnes KL, O'Grady JG, Hemer M, Macadam I, Abbs DJ, White CJ, Bennett JC, Corney SP, Holz GK, Grose MR, Gaynor SM & Bindoff NL In Press, Climate Futures for Tasmania: extreme tide and sea level events technical report, Antarctic Climate and Ecosystems Cooperative Research Centre, Hobart, Tasmania
- White CJ, Sanabria LA, Grose MR, Corney SP, Bennett JC, Holz GK, McInnes KL, Cechet RP, Gaynor SM & Bindoff NL 2011, Climate Futures for Tasmania: extreme events technical report, Antarctic Climate and Ecosystems Cooperative Research Centre, Hobart, Tasmania

The material in this report is based on computer modelling projections for climate change scenarios and, as such, there are inherent uncertainties involved. While every effort has been made to ensure the material in this report is accurate, Antarctic Climate & Ecosystems Cooperative Research Centre (ACE) provides no warranty, guarantee or representation that material is accurate, complete, up to date, non-infringing or fit for a particular purpose. The use of the material is entirely at the risk of a user. The user must independently verify the suitability of the material for its own use.

To the maximum extent permitted by law, ACE, its participating organisations and their officers, employees, contractors and agents exclude liability for any loss, damage, costs or expenses whether direct, indirect, consequential including loss of profits, opportunity and third party claims that may be caused through the use of, reliance upon, or interpretation of the material in this report.